

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Unintentional Injuries/Violence

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN8: Percentage of students who rarely or never wore a bicycle helmet (among students who had ridden a bicycle during the 12 months before the survey)	88.1 (84.2-91.2)	524	90.0 (83.3-94.1)	86			●
QN9: Percentage of students who rarely or never wore a seat belt (when riding in a car driven by someone else)	5.1 (3.6-7.1)	847	12.5 (6.9-21.7)	132		●	
QN10: Percentage of students who rode with a driver who had been drinking alcohol (in a car or other vehicle one or more times during the 30 days before the survey)	28.1 (24.8-31.7)	851	17.8 (12.5-24.6)	131	●		
QN11: Percentage of students who drove when drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	8.4 (4.3-15.7)	362	13.5 (7.2-24.0)	57			●
QN12: Percentage of students who texted or e-mailed while driving a car or other vehicle (on at least 1 day during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	26.5 (19.4-35.1)	375	29.8 (15.9-48.8)	59			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Unintentional Injuries/Violence

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN13: Percentage of students who carried a weapon (such as a gun, knife, or club on at least 1 day during the 30 days before the survey)	11.6 (9.2-14.7)	823	7.7 (4.2-13.5)	119			●
QN14: Percentage of students who carried a gun (on at least 1 day during the 30 days before the survey)	2.8 (2.0-3.9)	839	4.7 (1.9-11.2)	127			●
QN15: Percentage of students who carried a weapon on school property (such as a gun, knife, or club on at least 1 day during the 30 days before the survey)	4.6 (2.8-7.4)	841	4.0 (1.4-10.6)	123			●
QN16: Percentage of students who did not go to school because they felt unsafe at school or on their way to or from school (on at least 1 day during the 30 days before the survey)	7.8 (5.0-11.9)	847	15.1 (8.1-26.5)	128			●
QN17: Percentage of students who were threatened or injured with a weapon on school property (such as a gun, knife, or club one or more times during the 12 months before the survey)	6.6 (4.0-10.5)	855	9.8 (5.2-18.0)	133			●
QN18: Percentage of students who were in a physical fight (one or more times during the 12 months before the survey)	24.6 (20.9-28.7)	829	28.1 (18.0-41.1)	122			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Unintentional Injuries/Violence

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN19: Percentage of students who were injured in a physical fight (one or more times during the 12 months before the survey; injuries had to be treated by a doctor or nurse)	3.1 (1.9-5.0)	839	4.0 (1.6-9.5)	126			●
QN20: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)	9.5 (6.9-12.9)	836	13.3 (5.0-30.7)	126			●
QN21: Percentage of students who were ever physically forced to have sexual intercourse (when they did not want to)	9.4 (7.2-12.3)	843	15.2 (8.2-26.6)	126			●
QN22: Percentage of students who experienced physical dating violence (one or more times during the 12 months before the survey, including being hit, slammed into something, or injured with an object or weapon on purpose by someone they were dating or going out with among students who dated or went out with someone during the 12 months before the survey)	8.1 (6.1-10.8)	654	21.4 (12.9-33.5)	80		●	
QN23: Percentage of students who experienced sexual dating violence (one or more times during the 12 months before the survey, including kissing, touching, or being physically forced to have sexual intercourse when they did not want to by someone they were dating or going out with among students who dated or went out with someone during the 12 months before the survey)	8.4 (6.0-11.5)	661	26.1 (19.8-33.6)	84		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Unintentional Injuries/Violence

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN24: Percentage of students who were bullied on school property (during the 12 months before the survey)	14.6 (11.6-18.0)	846	26.8 (16.5-40.4)	125			●
QN25: Percentage of students who were electronically bullied (including being bullied through e-mail, chat rooms, instant messaging, websites, or texting during the 12 months before the survey)	11.8 (9.2-15.1)	843	17.1 (9.5-28.9)	125			●
QN26: Percentage of students who felt sad or hopeless (almost every day for 2 or more weeks in a row so that they stopped doing some usual activities during the 12 months before the survey)	32.8 (27.2-38.8)	852	56.5 (47.1-65.5)	128		●	
QN27: Percentage of students who seriously considered attempting suicide (during the 12 months before the survey)	15.5 (13.0-18.4)	848	31.6 (23.4-41.1)	121		●	
QN28: Percentage of students who made a plan about how they would attempt suicide (during the 12 months before the survey)	15.7 (13.4-18.2)	849	33.5 (23.2-45.5)	123		●	
QN29: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)	8.8 (6.5-11.7)	722	19.8 (14.8-26.1)	102		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

Los Angeles High School Survey  
Risk Behaviors and Sex of Sexual Contacts Report

Unintentional Injuries/Violence

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>‡</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>‡</sup>	No Difference <sup>‡</sup>
	% 95% CI <sup>*</sup>	N	% 95% CI	N			
QN30: Percentage of students who attempted suicide that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	2.3 (1.3-4.1)	711	3.8 (1.5-9.0)	101			●

<sup>\*</sup>Confidence interval.

<sup>‡</sup>Based on t-test analyses, p<0.05.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Tobacco Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN31: Percentage of students who ever tried cigarette smoking (even one or two puffs)	36.0 (32.1-40.1)	821	49.3 (37.3-61.4)	121			●
QN32: Percentage of students who smoked a whole cigarette before age 13 years (for the first time)	4.6 (3.1-6.9)	809	12.3 (5.8-24.2)	119			●
QN33: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)	6.4 (4.5-9.2)	797	19.4 (10.7-32.5)	116		●	
QNFRCIG: Percentage of students who currently frequently smoked cigarettes (on 20 or more days during the 30 days before the survey)	1.1 (0.6-2.1)	797	6.0 (2.2-15.3)	116			●
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)	0.6 (0.4-0.9)	797	5.7 (2.0-15.2)	116			●
QN34: Percentage of students who smoked more than 10 cigarettes per day (during the 30 days before the survey among students who currently smoked cigarettes on the days they smoked)	0.5 (0.1-3.4)	62	-	23			

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

- = Fewer than 30 students in the denominator (subgroup).

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Tobacco Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN35: Percentage of students who usually obtained their own cigarettes by buying them in a store or gas station (during the 30 days before the survey among students who currently smoked cigarettes and who were aged <18 years)	24.2 (9.1-50.7)	37	-	19			
QNCIGINT: Percentage of students who usually obtained their own cigarettes by buying on the internet (during the 30 days before the survey among students who currently smoked cigarettes and who were aged <18 years)	0.0 (.-)	37	-	19			
QN36: The percentage of students who did not try to quit smoking cigarettes (among students who currently smoked cigarettes during the 12 months before the survey)	44.8 (31.0-59.5)	60	-	21			
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, or dip on at least 1 day during the 30 days before the survey)	2.7 (1.7-4.4)	855	11.9 (6.3-21.3)	131		●	
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars on at least 1 day during the 30 days before the survey)	5.3 (3.7-7.5)	855	15.2 (8.4-25.8)	132		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

- = Fewer than 30 students in the denominator (subgroup).

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

Los Angeles High School Survey  
Risk Behaviors and Sex of Sexual Contacts Report

Tobacco Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN39: Percentage of students who ever used electronic vapor products (e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens such as blu, NJOY, or Starbuzz)	60.4 (54.1-66.4)	831	59.4 (47.8-70.0)	125			●
QN40: Percentage of students who currently used electronic vapor products (e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens such as blu, NJOY, or Starbuzz on at least 1 day during the 30 days before the survey)	24.1 (20.2-28.6)	836	28.8 (20.1-39.4)	130			●
QNTOB4: Percentage of students who currently used tobacco (current cigarette, smokeless tobacco, cigar, or electronic vapor product use on at least 1 day during the 30 days before the survey)	28.8 (23.6-34.6)	817	43.2 (29.5-58.1)	129			●
QNTOB3: Percentage of students who currently used cigarettes, cigars, or smokeless tobacco (on at least 1 day during the 30 days before the survey)	9.9 (7.4-13.1)	820	28.3 (16.6-44.1)	126		●	
QNTOB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)	8.9 (6.6-12.0)	811	25.4 (14.0-41.6)	122		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.



## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Alcohol/Other Drug Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN41: Percentage of students who ever drank alcohol (at least one drink of alcohol on at least 1 day during their life)	71.5 (66.5-75.9)	802	76.6 (63.2-86.1)	122			●
QN42: Percentage of students who drank alcohol before age 13 years (for the first time other than a few sips)	19.1 (16.3-22.2)	851	31.2 (21.7-42.7)	131		●	
QN43: Percentage of students who currently drank alcohol (at least one drink of alcohol on at least 1 day during the 30 days before the survey)	35.3 (29.5-41.5)	743	41.7 (31.9-52.2)	108			●
QN44: Percentage of students who drank five or more drinks of alcohol in a row (within a couple of hours on at least 1 day during the 30 days before the survey)	16.9 (12.9-21.8)	801	25.5 (16.4-37.3)	120			●
QN45: Percentage of students who reported that the largest number of drinks they had in a row was 10 or more (within a couple of hours during the 30 days before the survey)	4.1 (2.5-6.7)	742	6.1 (2.8-12.7)	111			●
QN46: Percentage of students who usually obtained the alcohol they drank by someone giving it to them (among students who currently drank alcohol)	39.2 (31.8-47.0)	254	23.3 (13.8-36.5)	52	●		

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

Los Angeles High School Survey  
Risk Behaviors and Sex of Sexual Contacts Report

Alcohol/Other Drug Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN47: Percentage of students who ever used marijuana (one or more times during their life)	53.3 (48.4-58.2)	820	65.7 (57.4-73.2)	121		●	
QN48: Percentage of students who tried marijuana before age 13 years (for the first time)	12.2 (9.3-15.8)	842	17.6 (8.7-32.6)	122			●
QN49: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)	27.3 (23.9-30.9)	840	38.1 (28.2-49.2)	126			●
QN50: Percentage of students who ever used cocaine (any form of cocaine, such as powder, crack, or freebase, one or more times during their life)	8.4 (6.3-11.1)	829	17.7 (9.8-30.1)	132			●
QN51: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	9.1 (6.4-12.9)	849	26.8 (17.8-38.3)	132		●	
QN52: Percentage of students who ever used heroin (also called "smack," "junk," or "China white," one or more times during their life)	1.7 (0.9-3.3)	847	10.0 (5.0-18.9)	132		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Alcohol/Other Drug Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN53: Percentage of students who ever used methamphetamines (also called "speed," "crystal," "crank," or "ice," one or more times during their life)	4.6 (2.8-7.6)	842	12.5 (6.7-22.3)	133		●	
QN54: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)	7.1 (5.3-9.6)	849	18.6 (11.2-29.4)	132		●	
QN55: Percentage of students who ever used synthetic marijuana (also called "K2", "Spice", "fake weed", "King Kong", "Yucatan Fire", "Skunk", or "Moon Rocks", one or more times during their life)	10.1 (8.4-12.1)	849	18.8 (12.3-27.7)	131		●	
QN56: Percentage of students who ever took steroids without a doctor's prescription (pills or shots, one or more times during their life)	2.9 (2.0-4.3)	851	10.0 (5.2-18.2)	131		●	
QN57: Percentage of students who ever took prescription drugs without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, one or more times during their life)	14.7 (12.0-17.7)	850	31.2 (20.5-44.2)	132		●	
QN58: Percentage of students who ever injected any illegal drug (used a needle to inject any illegal drug into their body one or more times during their life)	2.2 (1.2-3.8)	851	6.5 (3.0-13.4)	124			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

Los Angeles High School Survey  
Risk Behaviors and Sex of Sexual Contacts Report

Alcohol/Other Drug Use

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI <sup>*</sup>	N	% 95% CI	N			
QN59: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)	35.9 (29.9-42.4)	844	42.6 (31.2-54.9)	124			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Sexual Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN60: Percentage of students who ever had sexual intercourse	67.9 (61.5-73.7)	725	49.5 (37.7-61.2)	106	●		
QN61: Percentage of students who had sexual intercourse before age 13 years (for the first time)	8.1 (6.0-10.8)	732	11.0 (5.8-19.9)	109			●
QN62: Percentage of students who had sexual intercourse with four or more persons (during their life)	13.5 (10.6-17.1)	736	11.2 (5.4-21.8)	105			●
QN63: Percentage of students who were currently sexually active (sexual intercourse with at least one person during the 3 months before the survey)	46.7 (41.1-52.4)	732	32.0 (21.2-45.2)	104	●		
QN64: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	15.8 (12.7-19.7)	350	32.7 (20.1-48.4)	35		●	
QN65: The percentage of students who did not use a condom (during last sexual intercourse among students who were currently sexually active)	36.7 (30.4-43.5)	340	50.9 (29.0-72.4)	34			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Sexual Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN66: The percentage of students who did not use birth control pills (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	93.6 (89.8-96.1)	342	82.6 (57.4-94.4)	35			●
QNIUDIMP: Percentage of students who did not use an IUD (e.g., Mirena or ParaGard) or implant (e.g., Implanon or Nexplanon) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	97.3 (93.6-98.9)	342	90.3 (63.3-98.1)	35			●
QNSHPARG: Percentage of students who did not use a shot (e.g., Depo-Provera), patch (e.g., OrthoEvra), or birth control ring (e.g., NuvaRing) (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	95.0 (91.4-97.2)	342	93.9 (77.0-98.6)	35			●
QNOTHHPL: Percentage of students who did not use birth control pills; an IUD or implant; or a shot, patch, or birth control ring (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	86.0 (79.3-90.7)	342	66.9 (47.5-81.9)	35	●		
QNDUALBC: Percentage of students who did not use both a condom during and birth control pills; an IUD or implant; or a shot, patch, or birth control ring before last sexual intercourse (to prevent STD and pregnancy among students who were currently sexually active)	96.9 (94.2-98.4)	337	90.3 (72.6-97.0)	34			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

Los Angeles High School Survey  
Risk Behaviors and Sex of Sexual Contacts Report

Sexual Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI <sup>*</sup>	N	% 95% CI	N			
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy (during last sexual intercourse among students who were currently sexually active)	15.4 (10.8-21.4)	342	21.7 (8.1-46.5)	35			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Dietary Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>‡</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>‡</sup>	No Difference <sup>‡</sup>
	% 95% CI*	N	% 95% CI	N			
QNOBES: Percentage of students who were obese ( $\geq$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth chart <sup>‡</sup> )	10.1 (8.6-11.8)	767	19.2 (11.6-30.1)	113			●
QNOWT: Percentage of students who were overweight ( $\geq$ 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth chart <sup>‡</sup> )	19.3 (16.2-22.9)	767	28.4 (18.1-41.6)	113			●
QN69: Percentage of students who described themselves as slightly or very overweight	31.4 (27.4-35.7)	834	49.8 (39.6-60.0)	129		●	
QN70: The percentage of students who were not trying to lose weight	48.8 (45.0-52.6)	847	33.8 (26.2-42.3)	129	●		
QN71: Percentage of students who did not drink fruit juice (100% fruit juices one or more times during the 7 days before the survey)	19.4 (15.0-24.8)	849	23.2 (11.4-41.4)	127			●
QN72: Percentage of students who did not eat fruit (one or more times during the 7 days before the survey)	9.0 (6.9-11.7)	845	18.0 (11.1-27.7)	128		●	

\*Confidence interval.

<sup>‡</sup>Based on t-test analyses,  $p < 0.05$ .

<sup>‡</sup>Based on reference data from the 2000 CDC Growth Charts.



## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Dietary Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QNFR0: Percentage of students who did not eat fruit or drink 100% fruit juices (during the 7 days before the survey)	3.7 (2.9-4.9)	844	9.2 (4.1-19.1)	126			●
QNFR1: Percentage of students who did not eat fruit or drank 100% fruit juices one or more times per day (during the 7 days before the survey)	35.3 (32.2-38.5)	844	39.3 (26.3-54.0)	126			●
QNFR2: Percentage of students who did not eat fruit or drank 100% fruit juices two or more times per day (during the 7 days before the survey)	63.2 (60.1-66.1)	844	72.2 (62.9-79.9)	126		●	
QNFR3: Percentage of students who did not eat fruit or drank 100% fruit juices three or more times per day (during the 7 days before the survey)	75.7 (72.1-78.9)	844	82.0 (73.1-88.4)	126			●
QN73: Percentage of students who did not eat salad (during the 7 days before the survey)	42.8 (38.6-47.0)	848	33.7 (22.8-46.8)	129			●
QN74: Percentage of students who did not eat potatoes (during the 7 days before the survey)	35.1 (30.6-39.8)	843	30.5 (23.8-38.1)	129			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Dietary Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN75: Percentage of students who did not eat carrots (during the 7 days before the survey)	46.2 (43.0-49.5)	839	39.0 (25.0-55.0)	128			●
QN76: Percentage of students who did not eat other vegetables (during the 7 days before the survey)	20.4 (17.4-23.8)	846	22.8 (14.2-34.4)	128			●
QNVEG0: Percentage of students who did not eat vegetables (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	8.0 (5.7-11.0)	833	4.6 (1.6-12.4)	128			●
QNVEG1: Percentage of students who did not eat vegetables one or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	45.4 (40.4-50.5)	833	32.6 (21.3-46.4)	128			●
QNVEG2: Percentage of students who did not eat vegetables two or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	77.0 (73.0-80.6)	833	68.4 (59.3-76.3)	128			●
QNVEG3: Percentage of students who did not eat vegetables three or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	87.9 (84.2-90.8)	833	82.0 (74.1-87.9)	128			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Dietary Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN77: The percentage of students who drank a can, bottle, or glass of soda or pop (not including diet soda or diet pop, during the 7 days before the survey)	75.4 (70.7-79.5)	843	74.5 (65.8-81.5)	128			●
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	13.9 (10.9-17.5)	843	16.7 (9.2-28.5)	128			●
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	7.0 (5.3-9.2)	843	8.9 (3.9-19.1)	128			●
QNSODA3: Percentage of students who drank a can, bottle, or glass of soda or pop three or more times per day (not including diet soda or diet pop, during the 7 days before the survey)	3.8 (2.8-5.1)	843	5.2 (2.1-12.5)	128			●
QN78: Percentage of students who did not drink milk (during the 7 days before the survey)	19.8 (17.3-22.6)	831	25.1 (14.4-40.1)	129			●
QNMILK1: Percentage of students who did not drink one or more glasses per day of milk (during the 7 days before the survey)	68.5 (63.2-73.4)	831	75.3 (64.7-83.5)	129			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

Los Angeles High School Survey  
Risk Behaviors and Sex of Sexual Contacts Report

Dietary Behaviors

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QNMILK2: Percentage of students who did not drink two or more glasses per day of milk (during the 7 days before the survey)	80.5 (75.3-84.8)	831	87.3 (79.4-92.5)	129			●
QNMILK3: Percentage of students who did not drink three or more glasses per day of milk (during the 7 days before the survey)	92.3 (89.2-94.6)	831	94.8 (89.3-97.6)	129			●
QN79: Percentage of students who did not eat breakfast (during the 7 days before the survey)	11.2 (9.4-13.3)	840	21.4 (15.2-29.2)	128		●	
QNBK7DAY: Percentage of students who did not eat breakfast on all 7 days (during the 7 days before the survey)	64.8 (61.3-68.2)	840	82.5 (74.0-88.6)	128		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

Los Angeles High School Survey  
Risk Behaviors and Sex of Sexual Contacts Report

Physical Activity

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN80: The percentage of students who were not physically active at least 60 minutes per day on 5 or more days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	50.5 (45.7-55.3)	834	64.8 (51.9-75.8)	128		●	
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	13.6 (10.4-17.6)	834	19.6 (12.5-29.5)	128			●
QNPA7DAY: Percentage of students who were not physically active at least 60 minutes per day on all 7 days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	72.5 (67.4-77.1)	834	82.1 (73.0-88.6)	128		●	
QN81: Percentage of students who watched television 3 or more hours per day (on an average school day)	27.4 (24.0-31.1)	823	29.9 (20.8-41.0)	126			●
QN82: Percentage of students who played video or computer games or used a computer 3 or more hours per day (for something that was not school work on an average school day)	42.9 (39.1-46.8)	833	44.0 (37.3-51.0)	127			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Physical Activity

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN83: The percentage of students who did not attend physical education classes on 1 or more days (in an average week when they were in school)	47.2 (38.6-56.0)	825	49.5 (35.6-63.5)	126			●
QNDLYPE: Percentage of students who did not attend physical education classes on all 5 days (in an average week when they were in school)	71.4 (61.3-79.7)	825	75.5 (57.0-87.7)	126			●
QN84: The percentage of students who did not play on at least one sports team (run by their school or community groups during the 12 months before the survey)	43.1 (37.7-48.8)	831	51.4 (38.3-64.3)	127			●
QN88: The percentage of students who did not have 8 or more hours of sleep (on an average school night)	71.1 (66.0-75.6)	827	77.4 (65.9-85.8)	127			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

Los Angeles High School Survey  
Risk Behaviors and Sex of Sexual Contacts Report

Other	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	Health Risk Behavior	% 95% CI*	N	% 95% CI			
	QN85: The percentage of students who were never tested for HIV (not including tests done when donating blood)	79.5 (75.9-82.6)	830	83.3 (75.3-89.2)	126		●
	QN86: The percentage of students who did not see a dentist (for a check-up, exam, teeth cleaning, or other dental work during the 12 months before the survey)	29.8 (24.3-35.9)	832	33.1 (25.3-41.9)	126		●
	QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma	22.6 (19.3-26.4)	830	24.6 (16.5-34.9)	124		●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.

## 2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

### Los Angeles High School Survey Risk Behaviors and Sex of Sexual Contacts Report

#### Site-Added

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI*	N	% 95% CI	N			
QN90: Percentage of students who have been the victim of teasing or name calling because someone thought they were gay, lesbian, or bisexual (during the 12 months before the survey)	8.3 (6.4-10.6)	853	31.7 (22.0-43.3)	126		●	
QN91: Percentage of students who smoked cigarettes on school property (on at least 1 day during the 30 days before the survey)	2.0 (1.0-4.0)	849	9.7 (5.2-17.5)	131		●	
QN92: Percentage of students who have taken over-the-counter drugs to get high (one or more times during their life)	8.4 (6.5-10.7)	854	20.7 (11.8-33.5)	131		●	
QN93: Percentage of students who had oral sex	58.4 (52.8-63.7)	837	54.1 (44.6-63.3)	125			●
QN94: Percentage of students who had anal sex	18.6 (15.2-22.6)	848	27.4 (18.0-39.4)	125			●
QN97: Percentage of students who went without eating for 24 hours or more to lose weight or to keep from gaining weight (also called fasting, during the 30 days before the survey)	12.2 (9.2-16.1)	835	22.8 (15.1-33.1)	119		●	
QN98: Percentage of students who took some diet pills, powders, or liquids without a doctor's advice to lose weight or to keep from gaining weight (during the 30 days before the survey)	7.0 (5.3-9.2)	844	19.1 (12.7-27.9)	123		●	

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.



2015 YOUTH RISK BEHAVIOR SURVEY RESULTS

Los Angeles High School Survey  
Risk Behaviors and Sex of Sexual Contacts Report

Site-Added

Health Risk Behavior	Opposite Sex Only		Same Sex Only or Both Sexes		Opposite Sex Only More Likely Than Same Sex Only or Both Sexes <sup>†</sup>	Same Sex Only or Both Sexes More Likely Than Opposite Sex Only <sup>†</sup>	No Difference <sup>†</sup>
	% 95% CI <sup>*</sup>	N	% 95% CI	N			
QN99: Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight (during the 30 days before the survey)	4.9 (3.5-6.8)	840	21.6 (13.9-31.8)	121		●	
QN100: Percentage of students who have not been taught or are not sure if they have been taught about AIDS or HIV infection in school	21.0 (16.2-26.8)	831	23.1 (16.1-32.1)	126			●

\*Confidence interval.

<sup>†</sup>Based on t-test analyses, p<0.05.